

# High rate of ceftaroline-resistance in *Staphylococcus haemolyticus* causing late-onset sepsis and colonizing preterm neonates and their mothers

Hiie Soeorg<sup>1</sup>, Evamaria Elisabet Keränen<sup>1</sup>, Hanna Kadri Metsvaht<sup>1</sup>, Sirli Treumuth<sup>1</sup>, Imbi Eelmäe<sup>2</sup>, Mirjam Merila<sup>3</sup>, Mari-Liis Ilmoja<sup>4</sup>, Tuuli Metsvaht<sup>2</sup>, Irja Lutsar<sup>1</sup>

<sup>1</sup>Department of Microbiology, University of Tartu, Tartu, Estonia. <sup>2</sup>Pediatric Intensive Care Unit, Tartu University Hospital, Tartu, Estonia

<sup>3</sup>Department of Neonatology, Children's Clinic, Tartu University Hospital, Tartu, Estonia. <sup>4</sup>Pediatric Intensive Care Unit, Tallinn Children's Hospital, Tallinn, Estonia

## Background & Aim

- Ceftaroline is a new cephalosporin active against MRSA
- Susceptibility of *S. epidermidis* and *S. haemolyticus* causing late-onset sepsis (LOS) in preterm neonates warrants characterization before use of ceftaroline in treatment of LOS
- We aimed to determine susceptibility to ceftaroline in *mecA*-positive *S. epidermidis* and *S. haemolyticus* causing LOS and colonizing potential reservoirs of invasive strains (gut/skin of preterm neonates, mother's own milk (MOM))

## Materials and Methods

- A total of 97 previously characterized *mecA*-positive strains of diverse multilocus sequence types (Table 1)

Table 1. Number of isolates included in the study according to time, source and species

		Colonizing		Causing
		Gut/skin (n=41)	MOM (n=39)	LOS (n=17)
2007-	<i>S. epidermidis</i>	11	0	2
2008	<i>S. haemolyticus</i>	0	0	4
2014-	<i>S. epidermidis</i>	3	31	3
2015	<i>S. haemolyticus</i>	27	8	8

- Oxacillin and ceftaroline MICs were determined by MIC Test Strips (Liofilchem)
- Results interpreted according to EUCAST 2018 (Table 2)

Table 2. MIC breakpoints for determination of resistance (according to EUCAST 2018)

	Susceptible	Intermediate	Resistant
Ceftaroline*	≤1 mg/L	1<...≤2 mg/L	>2 mg/L
Oxacillin	≤0.25 mg/L	-	>0.25 mg/L

\*Ceftaroline breakpoint for *S. aureus* (indications other than pneumonia).

## Results

- *S. epidermidis* compared with *S. haemolyticus* had lower ceftaroline MICs ( $p < 0.001$ ) and lower resistance-rate to ceftaroline ( $p < 0.001$ ) (Table 3)

Table 3. The results of susceptibility testing to oxacillin and ceftaroline

		MIC <sub>50</sub>	MIC <sub>90</sub>	Susceptible (%)	Intermediate (%)	Resistant (%)
Oxacillin	<i>S. epidermidis</i>	2	16	4	0	96
	<i>S. haemolyticus</i>	≥256	≥256	0	0	100
Ceftaroline	<i>S. epidermidis</i>	0.25	0.5	98	0	2
	<i>S. haemolyticus</i>	3	4	6.4	23.4	70.2

- *S. epidermidis* from MOM had lower ceftaroline MICs compared with *S. epidermidis* colonizing gut/skin of neonates or causing LOS (MIC<sub>50</sub>/MIC<sub>90</sub> 0.25/0.5 vs 0.38/0.5 mg/L;  $p = 0.012$  and  $p = 0.02$ , respectively), but colonizing and LOS-causing *S. haemolyticus* had similar MICs (Figure 1)
- Ceftaroline-resistant *S. haemolyticus* isolates belonged to various sequence types (ST1, 3, 25, 42) and were detected in both time periods (2007-2008 and 2014-2015)




Figure 1. Distribution of MICs of ceftaroline in *S. epidermidis* and *S. haemolyticus* causing LOS or colonizing gut or skin of preterm neonates or MOM

## Conclusion

- *S. epidermidis* colonizing gut/skin and MOM or causing LOS in preterm neonates is mostly susceptible to ceftaroline.
- Widespread ceftaroline-resistance in *S. haemolyticus* may limit its potential use in treatment of LOS in preterm neonates.

### Acknowledgments:

The study was supported by the Target Financing of Estonian Ministry of Education and Research (SF0180004s12), European Regional Development Fund (project SFOS WP1-NeuroAIDS), Archimedes Foundation (Project No. 3.2.1001.11-0032), the European Society for Paediatric Infectious Diseases (ESPID Small Grant Award), Estonian Ministry of Education and Research (Grant No. KOGU-HUMB). Presenting author contact: hiie.soeorg@ut.ee