



EQUANIMITY®

IDENTIFICATION

Name: Equanimity®
Composition: Synthetic analog of maternal equine
Semiocchemical: 1%.
Excipient q.s. 5 ml.
Form: Single 5 ml stick of gel (1 stick = 1 application to spread at both nostrils).
be Licensee/Operator:
Semiokeys AW

Licence Holder: IRSEA.
Patents: n°98400117.2; n°99903663.5; n°2227706;
n°980298; n°2318392; n° PI 9907151-7;
n°99/0426; n°980282; n°09/010.515; n°09/525.763;
n°09/621.554; n°2000; n°P 341 867; n°505813;
n°137257; n°PO101312; n°99803663.5; n°24237/99;
n°2000-528279; n°007189; n°P990100248;
Manufacturer: ACP, Zone Industrielle de l'Europe —
BP59— 49260 Montreuil-Bellay — France.

PROPERTIES

Equanimity® reduces equine stress levels in horses allowing the animal to focus on its task.
— for prevention purposes, Equanimity® can be used alone.
— Equanimity® is recommended as part of reeducation programs. For optimal results, Equanimity® should be administered in a quiet place prior to the reeducation exercises.

WHEN TO USE Equanimity®

Equanimity® is recommended for young horses, for all horses exposed to a new situation or exercise, and for horses that are nervous, highly-sensitive to any new feature in their environment, as follows:
— transport (truck, airplane, train): boarding, journey, arrival;
— change of environment: movement, outdoor exercise, stable change, new owner;
— working conditions: breaking in a foal, introduction of a new exercise;
— social conditions: separation of the foal from the mare, time alone, meeting other horses;
— care and treatment: visit to the vet, blacksmith farrier, grooming;
— noise and light: storm, explosion, fireworks, crowds, motorized vehicles, flashing lights.

INSTRUCTIONS FOR USE

Rub the stick for 30 to 40 seconds.
Administer Equanimity® in a quiet place 20 to 30 minutes prior to the stress-inducing situation.
Place half a dose, 2.5 ml, on your finger and apply the gel to the base of the nostril; repeat the operation with the other half of the dose for the other nostril.
(5ml=quantity of gel required to reassure the horse for 2 to 2 ½ hours).
Repeat every 2 to ½ hours as required.

WARNING

Avoid contact with eyes. In case of contact with eyes, rinse thoroughly with water.
Keep out of reach of children.
Store at room temperature and away from sunlight.

CATEGORY

Health and well-being product.

NB:
Equanimity® is neither a painkiller nor an anesthetic and should not be used as such.
In the event of ongoing behavioral disorder, a complete clinical examination by a veterinarian is recommended to establish a diagnosis and suitable therapy program.



| Semiokeys

Semiokeys Animal Word
Quartier Saligman
84400 Apt - France
Tel: +33 (0) 4 90 75 06 04
www.semiokeys.com
www.equanimity.pro